



Contents lists available at ScienceDirect

Schizophrenia Research

journal homepage: www.elsevier.com/locate/schres

Letter to Editors

Context-sensitive social cognition is impaired in schizophrenic patients and their healthy relatives

Social cognition performance has been extensively studied within the schizophrenic population since Frith (1992) proposed a model relating mentalizing deficits and symptoms of the disorder. Moreover, recent research has suggested that social cognition is also impaired in unaffected relatives. Their deficits seem to be more pronounced when highly sophisticated tests are used (Janssen et al., 2003).

Fifteen clinically-stable medicated schizophrenic patients from multiplex families, fourteen healthy first-degree relatives, and eighteen controls were examined. Groups were similar in age, education, and sex distribution.

Initial selection criteria for all patients were: (1) diagnosis of paranoid schizophrenia according to DSM-IV-TR criteria (APA, 2000) and confirmed with Schedules of Clinical Assessment in Neuropsychiatry (SCAN) applied to both the patients and their relatives; and (2) the presence of one or more relatives with the diagnosis of schizophrenia (no greater than third-degree relative), evidenced by the Family Interview for Genetic Studies (FIGS). Healthy relatives had to be first-degree relatives. SCAN was applied to relatives in order to rule out any psychiatric conditions in this group. All participants were between 20 and 55 years old. Participants completed written informed consent and were paid for their participation in the study.

Participants completed the Raven's Coloured Progressive Matrices, Trail Making Test, Reading the Mind in the Eyes, and Faux Pas Test. Additionally, Positive and Negative Syndrome Scale (PANSS) and Calgary Depression Scale for Schizophrenia were applied to patients to characterize symptomatology and mood. Detailed information about participants, results and discussion is presented as [Supplementary data](#). Main results (Table 1) were:

- Raven's Progressive Matrices: Similar intellectual functioning was observed between the groups ($p > 0.05$)
- Trail Making Test-A: No difference was found ($p > 0.05$)
- Trail Making Test-B: Significant difference was found ($p < 0.05$). However, post hoc comparison only showed differences between patients and controls ($p < 0.05$)
- Reading the Mind in the Eyes: Similar performance in emotion recognition was observed across the groups ($p > 0.05$)
- Faux Pas: A significant difference between groups was found ($p < 0.001$). Post hoc analysis showed worse performance of patients and relatives compared to controls ($p < 0.001$ and

$p < 0.05$, respectively). No differences between patients and relatives were found ($p > 0.05$)

The Faux Pas impairment in patients replicated data reported elsewhere (Shur et al., 2008). However, to the best of our knowledge, there are no previous data reporting that Faux Pas performance is impaired in healthy relatives.

Unexpectedly, no significant differences between the groups were found in the Mind in the Eyes Test. Despite several studies reporting impaired performance on this test, other studies have reported otherwise (Kelemen et al., 2004; Kington et al., 2000).

The Faux Pas finding does not seem to be explained by IQ level, since the three groups performed similarly on the Raven's test. Instead, the Faux Pas deficit observed in patients and relatives suggests that it might be due to impaired processing of context-sensitive information. However, performance on the Faux Pas may be partially influenced by working memory load of the task and executive functioning. Future studies should address this issue in larger samples.

Social cognition is a collection of inferential abilities achieved during neurodevelopment, and the ability to detect a faux pas achieved later in development relative to other social cognition skills (Stone et al., 1998). Detecting a faux pas requires the integration of various components, namely (1) that the person committing faux pas does not know that he/she should not say it; and (2) the empathic understanding of the listener's feelings. The Faux Pas test requires integrating cognitive and emotional information in a context-sensitive manner to regulate behavior. For that reason, the Faux Pas test may be more sensitive in detecting deficits in healthy relatives.

Considering the limitations of this study, our results suggest that the high-level components of social cognition may be considered as possible endophenotype candidates for future studies.

Role of Funding Source

Funding for this study was provided by CONICET Grant and for the FONDECYT and Diego Portales Grants; those institutions have no further role in study design; in the collection, analysis and interpretation of data; in the writing of the report; and in the decision to submit the paper for publication.

Appendix A. Supplementary data

Supplementary data associated with this article can be found in the online version, at [doi:10.1016/j.schres.2009.10.017](https://doi.org/10.1016/j.schres.2009.10.017).

Table 1

Performance for patients with schizophrenia, first-degree relatives, and controls.

	Schizophrenia (n = 15)	Relatives (n = 14)	Controls (n = 18)	ANOVA F value (df.) and significant post-hoc comparisons
Raven's Coloured Progressive Matrices	29.6(4.03)	29.21(6.94)	32(3.09)	F(2,45) = 1.51
Trail Making Test-A	55.0(24.86)	41.5(20.06)	43.57(21.9)	F(2,45) = 2.26
Trail Making Test-B	141.86(88.4)	94.7(55.1)	85.0(26.9)	F(2,45) = 4.3338* Patients–Controls*
Reading the Mind in the Eyes Test	24.53(6.15)	26.36(7.22)	27(4.16)	F(2,45) = 0.7709
Faux pas Test	24.8 (14.97)	30,07(10.43)	41.72(9.69)	F(2,45) = 9.2623*** Relatives–Controls* Patients–Controls***
PANNS				
Positive	17.73(6.06)			
Negative	18.67(6.2)			
General	28.6(10.4)			
Calgary Depression Scale for Schizophrenia	3.39(2.8)			

Mean scores and standard deviations (presented in parentheses). Tukey's post-hoc analysis was carried out where appropriate.

Mean (\pm SD); ** $P < 0.01$; *** $P < 0.001$.**References**

- American Psychiatric Association, 2000. Diagnostic and Statistical Manual, Fourth Edition, Text Revised. American Psychiatry Association, Washington.
- Frith, C.D., 1992. The Cognitive Neuropsychology of Schizophrenia. Psychology Press.
- Janssen, I., Krabbendam, L., Jolles, J., van Os, J., 2003. Alterations in theory of mind in patients with schizophrenia and non-psychotic relatives. *Acta Psychiatr. Scand.* 108, 110–117.
- Kelemen, O., Kéri, S., Must, A., Benedek, G., Janka, Z., 2004. No evidence for impaired 'theory of mind' in unaffected first-degree relatives of schizophrenia patients. *Acta Psychiatr. Scand.* 110, 146–149.
- Shur, S., Shamay-Tsoory, S., Levkovitz, Y., 2008. Integration of emotional and cognitive aspects of theory of mind in schizophrenia and its relation to prefrontal neurocognitive performance. *Cogn. Neuropsychiatry* 13 (6), 472–490.
- Stone, V., Baron-Cohen, S., Knight, R., 1998. Frontal lobe contributions to theory of mind. *J. Cogn. Neurosci.* 10, 640–656.
- Kington, J., Jones, L., Watt, A., Hopkin, E., Williams, J., 2000. Impaired eye expression recognition in schizophrenia. *J. Psychiatr. Res.* 34, 341–347.

R. Riveros
E. Hurtado
M. Escobar
A. Ibañez*

*Laboratory of Cognitive Neuroscience,
Universidad Diego Portales, Santiago, Chile*

*Corresponding author. Laboratory of Experimental Psychology & Neuroscience Institute of Cognitive Neurology (INECO) & CONICET Castex 3293 (CP 1425) Buenos Aires, Argentina. Tel./fax: +54 11 4807 4748.

URL: <http://www.neurologiacognitiva.org/> (A. Ibañez).

E-mail address: aibanez@neurologiacognitiva.org (A. Ibañez).

A. Ibañez

*National Scientific and Technical Research Council (CONICET),
Buenos Aires, Argentina*

F. Manes

M. Cetkovich

A. Ibañez

*Laboratory of Experimental Psychology,
Institute of Cognitive and Behavioural Neurology (INECO),
Argentina
Institute of Neuroscience, Favaloro University, Argentina*

E. Hurtado

*School of Engineering,
Pontificia Universidad Católica de Chile, Chile*

M. Martin Reyes

*Unidad de Hospitalización Psiquiátrica. Hosp. Virgen del Camino,
Pamplona, Spain*

17 September 2009

Available online xxxx